

NAVAL OCEANOGRAPHIC OFFICE SPECIAL PUBLICATION 1400-NA 1

SURFACE CURRENTS





THE PARTY OF THE P

DTIC ELECTE JUN 1 3 1980




DISCLAIMER NOTICE

THIS DOCUMENT IS BEST QUALITY PRACTICABLE. THE COPY FURNISHED TO DTIC CONTAINED A SIGNIFICANT NUMBER OF PAGES WHICH DO NOT REPRODUCE LEGIBLY.

6 SURFACE CURRENTS.


GREENLAND SEA INCLUDING DENMARK ST



APPROVED FOR PUBLIC RELEASE; DISTRIBUTION UNLIMITED.

NAVAL OCEANOGRAPHIC OFFICE NSTL STATION, MISSISSIPPI 39522


25





REACE CURRENTS.

SEA INCLUDING DENMARK STRAIT .









VED FOR PUBLIC RELEASE; DISTRIBUTION UNLIMITED.

AVAL OCEANOGRAPHIC OFFICE STL STATION, MISSISSIPPI 39522

250450 L'11

ACKNOWLEDGMENTS

Messrs. Raymond J. Beauchesne* and William E. Boisvert made major contributions to this atlas.

*Mr. Beauchesne presently is employed by the Bureau of Naval Personnel.

FOREWORD

THIS ATLAS, ONE IN A SERIES OF 43 REGIONAL SURFACE CURRENT ATLAS IS PRODUCED TO FULFILL A NEED OF NAVY PLANNING STAFFS AND THE SCIENTIFIC A INDUSTRIAL COMMUNITIES FOR THE LATEST AVAILABLE OCEAN SURFACE CURRENT DATA THESE ATLASES ADD TO THE WEALTH OF NAUTICAL INFORMATION UPON WHICH OPERATIONAL PLANNING, NAVIGATIONAL SAFETY, AND SHIPPING ECONOMY DEPEND. RAPID PRODUCTION AND WIDE DISSEMINATION OF THIS ATLAS ARE MADE POSSIBLE BY THE LATEST COMPUTER TECHNIQUES.

THE CONSTANT IMPROVEMENT IN THE QUALITY OF SURFACE CURRENT DATA RECEIVED OVER THE YEARS IS MADE POSSIBLE LARGELY BY THE MORE THOROUGH REPORT VOLUNTARY OBSERVERS IN RECENT YEARS. THE DEFENSE MAPPING AGENCY, THE OCEANOGRAPHIC OFFICE, AND THE USER OF THE ATLASES RELY ON THE PERSONAL OBSERVATIONS OF THE MAN WHO HAS "BEEN THERE." MARINERS, IN REPORTING THEIR OBSERVATIONS, RENDER A SERVICE NOT ONLY TO THEMSELVES BUT ALSO TO ALL "WHO GO DOWN TO THE SEA IN SHIPS." WITH THE ADVENT OF NUCLEAR POWER, ELECTRONINAVIGATION AIDS, AND 300,000-TON SHIPS, UP-TO-DATE, RAPIDLY DISSEMINATED ENVIRONMENTAL AND NAVIGATIONAL INFORMATION HAS BECOME INCREASINGLY IMPORTA

JOHN R. McDONNELL Captain, U.S. Navy

Commander

FOREWORD

IN A SERIES OF 43 REGIONAL SURFACE CURRENT ATLASES, SED OF NAVY PLANNING STAFFS AND THE SCIENTIFIC AND THE LATEST AVAILABLE OCEAN SURFACE CURRENT DATA. EALTH OF NAUTICAL INFORMATION UPON WHICH OPERAVAL SAFETY, AND SHIPPING ECONOMY DEPEND. RAPID INATION OF THIS ATLAS ARE MADE POSSIBLE BY THE

ROVEMENT IN THE QUALITY OF SURFACE CURRENT DATA MADE POSSIBLE LARGELY BY THE MORE THOROUGH REPORTS RECENT YEARS. THE DEFENSE MAPPING AGENCY, THE THE USER OF THE ATLASES RELY ON THE PERSONAL OBHAS "BEEN THERE." MARINERS, IN REPORTING THEIR VICE NOT ONLY TO THEMSELVES BUT ALSO TO ALL "WHO S." WITH THE ADVENT OF NUCLEAR POWER, ELECTRONIC NOT ONLY TO THE POSSIBLY DISSEMINATED TOWAL INFORMATION HAS BECOME INCREASINGLY IMPORTANT.

JOHN R. McDOWNELL Captain, U.S. Navy Commander

ACCESSION	for	
NTIS DDC UNANNOUNC JUSTIFICATI	White So Buff So CED	-
BY DISTRIBUTION	ON/AVAHABIL	ITY CODES OF SPECIAL
A A	270	
<u></u>	<u> </u>	0


SURFACE CURRENT ATLASES

THIS SERIES OF COMPUTERIZED ATLASES REPLACES THE OLD HYDROGRAPHIC OFFICE ATLASES OF SURFACE CURRENTS (HOP 566, 568, 569, 570) WHICH WERE MANUALLY COMPILED FROM DATA OBTAINED DURING THE PERIOD 1903 - 1934. THESE NEW ATLASES CONFORM TO THE STANDARD NAVY OCEAN AREA AND REGION INDEX LIMITS SHOWN BELOW: e.g., NOO SP 1402-NP 10 COVERS NORTH PACIFIC REGION 10 EAST OF THE PHILIPPINES.

AS AMOUNTS OF NEW DATA

THESE GRAPHICS MAY AREAS AS THE NORTH SEA. CURRENTS ARE STRONGLY TI PREDICTABLE HOURLY CHANG

RECENT IMPROVEMENTS IN THE DATA FILE ASSURE THE INCLUSION OF THE LATEST, HIGH QUALITY SURFACE CURRENT DATA AVAILABLE. THE FILE NOW CONTAINS MORE THAN 4,200,000 OBSERVATIONS AND A GENERAL UPDATE OF THE FILE WILL BE MADE




CURRENT ATLASES



AS AMOUNTS OF NEW DATA WARRANT, MOST LIKELY EVERY 12 - 18 MONTHS.

THESE GRAPHICS MAY NOT BE TRULY REPRESENTATIVE OF THE ACTUAL FLOW IN SUCH AREAS AS THE NORTH SEA, PERSIAN GULF, GULF OF THAILAND, AND YELLOW SEA WHERE CURRENTS ARE STRONGLY TIDAL. FOR SUCH AREAS, OTHER SOURCES DESCRIBING PREDICTABLE HOURLY CHANGES OF TIDAL CURRENTS SHOULD BE CONSULTED.



Introduction

The Surface Current Data File, from which these atlases are derived, consists primarily of over four million ship set and drift observations. These data were collected by the Netherlands, Japan, Britain, France, and the United States. The file is supplemented by several thousand Geomagnetic Electrokinetograph (GEK) observations, mostly Japanese. The file spans the period from the early 1850's to the present. The earliest observations were collected by the Netherlands and Great Britain; those of the 1960's through the present are primarily United States data.


General Quality

The quality of this data file is considered high for this type of derived value. The data have been carefully screened for duplication; observations taken under adverse conditions (i.e. high winds and waves, time between observations greater than 12 hours) have been eliminated when warranted. Consideration was given to the reliability of the observer; doubtful shipboard computations of set and drift were edited; and observations with erroneous locations (mostly observations on land) have been eliminated. The accepted data are considered most useful when used collectively as in summaries where a number of observations show trends.

General Observation Technique

The set (direction) and drift (speed) are computed by the navigator from the difference between the dead reckoning (DR) position and the position determined by any type of navigational fix. The drift can be determined along any straight line track and includes all factors which cause changes in the DR position. When a fix is obtained, the current set (direction) is FROM the DR position TO the fix; the drift (speed) is equal to the distance in nautical miles between the DR and the fix, divided by the number of hours since the last fix. For successive observations, the TO POSITION of one observation becomes the FROM POSITION of the next observation.

Because the influence of current may vary along a ship's track, the MEAN POSITION of the track is assigned as the geographic location of the current observation. An example of a current computation is shown in the figure below.




EXAMPLE OF A SURFACE CURRENT (SHIP'S DRIFT) OBSERVATION


Data Presentation

فاستعلاه بالمالي والماليان


The following legend shows two types of surface current presentations by 1° quadrangle, type 1 with 12 or more observations and type 2 with fewer than 12 observations. Where there are 11 or fewer observations within a 1° quadrangle, the total number of observations is shown within the 90° quadrant containing the observations.




If there are 12 or more observations in by vector resultants as follows:



- Persistent Current 60 percent or more of all observations fall within a 45° sector of the 8-point compass.
 - (2) Prevailing Cur all observation 45° sectors.



(4) <u>Bizonal Flow</u> - Practically all observations are concentrated in opposite pairs of 45° sectors, and one pair contains at least 80 percent as many observations as the opposite pair. This generally indicates variability that occurs in zones of entrainment between opposing currents (see examples A and B, quadrangles 1, 2, and 3).



If there are 12 or more observations in a 1^{*} quadrangle, the surface current is depicted by vector resultants as follows:

narily by the hted anese. ervarough


The se hours)

d oblnated.


erence of includes current the hours ion

N of example

 Persistent Cu:rent - 60 percent or more of all observations fall within a 45° sector of the 8-point compass.




(2) Prevailing Current - 70 percent or more of all observations fall within two adjacent 45° sectors.




(3) Primary Current with Secondary Direction (a) Primary Current - 50 percent or more of all observations fall within three adjacent 45° sectors.


(b) Secondary Direction ~ 20 percent or more of all observations fall within a 45° sector, and the two resultant vector directions are separated by more than 90° of arc.



(4) <u>Bizonal Flow</u> - Practically all observations are concentrated in opposite pairs of 45° sectors, and one pair contains at least 80 percent as many observations as the opposite pair. This generally indicates variability that occurs in zones of entrainment between opposing currents (see examples A and B, quadrangles 1, 2, and 3).



(5) Variable Current - The 45° sector with most observations has less than 25 percent of all observations; direction is indeterminate.






irangle, Where

valosa.


100 OR


OBSERVATIONS.


, 0.6 KSOT




Mar










*




Ø


•



ن ان



أدردني





A

DISTRIBUTION LIST

NAVY

CINCPACFLT (02M) COMTHIRDFLT COMSEVENTHFLT COMSUBPAC COMNAVAIRPAC COMPATWINGSPAC PATWINGSPAC DET ADAK PATWING 1 COMNAVSURFPAC **DI RNAVOCEANMET** FLENUMEACEN FLEWEACEN GUAM FLEWEACEN PEARL NAVWEASERVFAC SAN DIEGO NAVWEASERVFAC YOKOSUKA NWSD ASHEVILLE NWSED ADAK NWSED AGANA NWSED ATSUGI NWSED KADENA NWSED MISAWA

OTHER GOVT.

NOAA/NODC NOAA/NCC

PRIVATE & UNIVERSITIES

FLORIDA ST. UNIV.
LOUISIANA ST. UNIV.
MASS. INST. OF TECH
ORE. ST. UNIV.
TEXAS A&M UNIV.
UNIV. OF MIAMI
UNIV. OF R.I.
UNIV. OF WASH.
SCRIPPS INST OF OCEANOGRAPHY
WOODS HOLE OCEANOGRAPHIC INST.

FOREIGN

HYDROGRAPHER/R.A.N.
DEPT. TRANSPORTATION/AUSTRALIA

READ INSTRUCTIONS
BEFORE COMPLETING FORM
3 RECIPIENT'S CATALOG NUMBER REPORT DOCUMENTATION PAGE AD-A085441 REPORT NUMBER NOO SP1400 - NA1 TITLE (and Subtitle)
SURFACE CURRENTS TYPE OF REPORT & PERIOD COVERED GREENLAND SEA INCLUDING DENMARK STRAIT PERFORMING ORG. REPORT NUMBER CONTRACT OR GRANT NUMBER(+) AUTHOR(s) Naval Oceanographic Office NSTL Station
Bay St. Louis, MS 39522
PERFORMING ORGANIZATION NAME AND ADDRESS PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS Naval Oceanographic Office REPORT DATE CONTROLLING OFFICE NAME AND ADDRESS January 1978 15. SECURITY CLASS. (of this report) MONITORING AGENCY NAME & ADDRESS(II different from Controlling Office) ISE. DECLASSIFICATION/DOWNGRADING DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited. DISTRIBUTION STATEMENT (of the obstract entered in Block 20, If different from Report) SUPPLEMENTARY NOTES KEY WORDS (Continue on reverse side if necessary and identify by block number) Greenland Sea Denmark Strait ABSTRACT (Continue on reverse side if necessary and identify by block ma This atlas, and the series of which it is a part, is computer generated and automatically plotted. It makes available to user the most recent surface current data collected and will be updated whenever sufficient amounts

of data are added to the data file. This and the other atlases are based on a vast quantity of data as compared to the previous manually-compiled

SECURITY CLASSIFICATION OF THIS PAGE (When Date Entered)

editions printed in the mid-thirties.

1 JAN 73 1473

EDITION OF 1 NOV 68 IS 0850LETE 5/N 0102-014-6601

URITY CLASSIFICATION OF THIS PAGE (When Date Entered)

SECURITY CLASSIFICATION OF THIS PAGE When Data Entered

20. Continued

The surface current information is based mainly on ship drift, whi is the difference between the dead reckining position and the position determined by any type of navigational fix. This difference describes the direction and speed of the current.

SECURITY CLASSIFICATION OF THIS PAGE(When Date Enter

ATEL